



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Reply to ‘Is single portal vein perfusion the best approach for machine preservation of liver grafts?’

Schlegel, Andrea ; Kron, Philipp ; de Oliveira, Michelle L ; Clavien, Pierre-Alain ; Dutkowski, Philipp

DOI: <https://doi.org/10.1016/j.jhep.2016.01.033>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-125995>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Schlegel, Andrea; Kron, Philipp; de Oliveira, Michelle L; Clavien, Pierre-Alain; Dutkowski, Philipp (2016). Reply to ‘Is single portal vein perfusion the best approach for machine preservation of liver grafts?’. *Journal of Hepatology*, 64(5):1195-1196.

DOI: <https://doi.org/10.1016/j.jhep.2016.01.033>

Reply to „ Is single portal vein perfusion the best approach for machine preservation of liver grafts? “

Andrea Schlegel, Philipp Kron, Michelle L. de Oliveira, Pierre-Alain Clavien*, Philipp Dutkowski

Department of Surgery & Transplantation
University Hospital Zurich
Swiss HPB and Transplant Center
Raemistr. 100
8091 Zurich, Switzerland

Word count: 469

Figures: 1

* Corresponding author:

Pierre-Alain Clavien
Department of Visceral Surgery and Transplantation
University Hospital Zurich
Swiss HPB and Transplant Center
Raemistrasse 100, CH-8091 Zurich
Phone: +41 44 255 2300
Fax: +41 44 255 4999
E-mail: clavien@access.uzh.ch

To the Editor:

We appreciate the interest and comments expressed in the letter by Isabel et al. concerning our recent findings on single portal vein hypothermic liver graft perfusion prior to transplantation (1).

The authors argue that single portal vein perfusion in the cold may not be sufficient for preservation of the extrahepatic bile duct of liver grafts, in contrast to dual HOPE (portal vein plus hepatic artery). This statement is based on the fact, that our images lack information on histological scoring of the perivascular biliary plexus. The authors claim also direct measurement of oxygen saturation and ATP analysis of the bile duct wall during single portal vein perfusion. In addition, they refer to publications from 2003 and 1999 on the effects of dual (hepatic artery and portal vein) perfusion on biliary secretion (2,3).

We agree that ATP measurements in bile duct tissue could in theory further support the presence of oxygen for aerobic metabolism in biliary epithelial during single portal vein perfusion. We showed, however, that the extrahepatic bile duct wall in both rodent and human DCD liver grafts is completely stained with a perfusate containing fluoresceine and also oxygen (60-80 kPa) (4). Based on this, our study demonstrated that HOPE through the portal vein transports perfusate, loaded with a high amount of oxygen, to the entire extrahepatic bile duct.

To underline viability of the extrahepatic Choledochus after reperfusion, we provide detailed histological images and scoring of human DCD bile ducts after portal vein HOPE treatment and transplantation. Supporting an intact biliary epithelium, we failed to document any mural necrosis, vascular injury, or even deep peribiliary gland injury, and the absence of inflammation (Figure 1) (5).

Under normothermic perfusion conditions, as stated in the studies by Foley et al, dual blood perfusion supply is essential for biliary function and epithelial viability (2, 3). The hypothermic perfusion setting, however, fundamentally differs from normothermic physiological conditions. During cold machine liver perfusion, oxygen consumption of any liver cells, including the extrahepatic bile duct, is dramatically reduced, and we convincingly reported that single portal vein perfusion with high oxygen tension enables sufficient liver ATP reload in the cold before graft implantation (6-8). To support our observation, recent human application of single

portal vein HOPE in extended DCD livers displayed a significant reduction of biliary complications with robust graft survival benefit, compared to matched unperfused DCD livers (9).

In summary, we believe that single portal vein HOPE is very effective. It is currently the simplest machine perfusion approach, with therefore high practicability and low associated costs. We agree that randomized trials remain needed to identify the best liver perfusion strategy, but, it is rather unlikely that additional arterial perfusion during HOPE in the cold would add more oxygen availability to the intra- and extrahepatic biliary tract, and thereby to be of any benefits.

References

1. Brüggerwirth IMA, Burlage LC, Porte RJ, Martins PN, Is single portal vein perfusion the best approach for machine preservation of liver grafts? JHepatol 2016
2. Foley DP, Vittimberga FJ, Quarfordt SH, Donohue SE, Traylor AN, MacPhee J, McLaughlin T, Ricciardi R, Callery MP, Meyers WC, Biliary secretion of extracorporeal porcine livers with single and dual vessel perfusion, Transplantation. 1999 Aug 15;68(3):362-8
3. Foley DP, Ricciardi R, Traylor AN, McLaughlin TJ, Donohue SE, Wheeler SM, Meyers WC, Quarfordt SH, Effect of hepatic artery flow on bile secretory function after cold ischemia, Am J Transplant. 2003 Feb;3(2):148-55
4. Schlegel A, Kron P, De Oliveira ML, Clavien PA, Dutkowski P. Is single portal vein approach sufficient for hypothermic machine perfusion of DCD liver grafts? J Hepatol. 2016 Jan;64(1):239-41
5. Op den Dries S, Westerkamp AC, Karimian N, Gouw AS, Bruinsma BG, Markmann JF, Lisman T, Yeh H, Uygün K, Martins PN, Porte RJ. Injury to peribiliary glands and vascular plexus before liver transplantation predicts formation of non-anastomotic biliary strictures. J Hepatol. 2014 Jun;60(6):1172-9.
6. Dutkowski P, Odermatt B, Heinrich T, Schönfeld S, Watzka M, Winkelbach V, Krysiak M, Junginger T, Hypothermic oscillating liver perfusion stimulates ATP synthesis prior to transplantation, J Surg Res. 1998 Dec;80(2):365-72
7. Dutkowski P, Graf R, Clavien PA, Rescue of the cold preserved rat liver by hypothermic oxygenated machine perfusion. Am J Transplant. 2006 May;6(5 Pt 1):903-12
8. Schlegel A, Graf R, Clavien PA, Dutkowski P. Hypothermic oxygenated perfusion (HOPE) protects from biliary injury in a rodent model of DCD liver transplantation. J Hepatol. 2013 Nov;59(5):984-91
9. Dutkowski P, Polak WG, Muiesan P, Schlegel A, Verhoeven CJ, Scalera I, DeOliveira ML, Kron P, Clavien PA. First Comparison of Hypothermic Oxygenated PERfusion Versus Static Cold Storage of Human Donation After Cardiac Death Liver Transplants: An International-matched Case Analysis. Ann Surg. 2015 Nov;262(5):764-71

Figure legend

Figure 1: Extrahepatic bile duct of human DCD liver graft after single portal vein HOPE and transplantation (* luminal surface). The mural stroma of the bile duct wall is healthy without necrosis (A, B) (#). Higher magnification displayed no loss of epithelial cells in periluminal peribiliary glands (arrows)(C,D), and also normal appearance of deep peribiliary glands (arrows)(E,F). The peribiliary vascular plexus shows normal vascular structure (G).